

Publikationen: Prof. Dr. Steffen Schröter, Chefarzt Unfall- und Wiederherstellungschirurgie

1. Ahmad SS, Schreiner AJ, Hirschmann MT, Schröter S, Dobele S, Ahrend MD, et al. (2019) Dynamic intraligamentary stabilization for ACL repair: a systematic review. *Knee Surg Sports Traumatol Arthrosc* 27:13-20
2. Ahrend M, Ateschrang A, Dobele S, Stockle U, Grunwald L, Schröter S, et al. (2016) [Return to sport after surgical treatment of a posterior cruciate ligament injury : A retrospective study of 60 patients]. *Orthopade* 45:1027-1038
3. Ahrend M, Ateschrang A, Stockle U, Schröter S(2016) [Treatment strategy for posttraumatic complex deformity. After bilateral femoral shaft fractures]. *Unfallchirurg* 119:151-158
4. Ahrend MD, Schröter S(2020) Critical comments to the publication „Hevesi M, Macalena JA, Wu IT, Camp CL, Levy BA, Arendt EA, et al. (2018) High tibial osteotomy with modern PEEK implants is safe and leads to lower hardware removal rates when compared to conventional metal fixation: a multi-center comparison study. *Knee Surgery, Sports Traumatology, Arthroscopy* 1-11“. *Knee Surg Sports Traumatol Arthrosc* 28:662-663
5. Ateschrang A, Ahmad SS, Stockle U, Schröter S, Schenk W, Ahrend MD (2018) Recovery of ACL function after dynamic intraligamentary stabilization is resultant to restoration of ACL integrity and scar tissue formation. *Knee Surg Sports Traumatol Arthrosc* 26:589-595
6. Ateschrang A, Albrecht D, Schröter S, Weise K, Dolderer J (2011) Current concepts review: septic arthritis of the knee pathophysiology, diagnostics, and therapy. *Wien Klin Wochenschr* 123:191-197
7. Ateschrang A, Albrecht D, Schröter S, Hirt B, Weise K, Dolderer JH (2011) Septic arthritis of the knee: presentation of a novel irrigation-suction system tested in a cadaver study. *BMC Musculoskelet Disord* 12:180
8. Ateschrang A, Dobele S, Freude T, Stockle U, Schröter S, Kraus TM (2016) Acute MCL and ACL injuries: first results of minimal-invasive MCL ligament bracing with combined ACL single-bundle reconstruction. *Arch Orthop Trauma Surg* 136:1265-1272
9. Ateschrang A, Eggensperger F, Ahrend MD, Schröter S, Stockle U, Kraus TM (2018) Obesity causes poorer clinical results and higher re-tear rates in rotator cuff repair. *Arch Orthop Trauma Surg* 138:835-842
10. Ateschrang A, Fiedler S, Schröter S, Stockle U, Freude T, Kraus TM (2014) [Duration of inability for work and return to physical work after arthroscopic and open labrum refixation]. *Z Orthop Unfall* 152:252-259
11. Ateschrang A, Freude T, Grunwald L, Schaffler A, Stockle U, Schröter S(2014) [Patella dislocation: an algorithm for diagnostic and treatment considering the rotation]. *Z Orthop Unfall* 152:59-67
12. Ateschrang A, Korner D, Joisten K, Ahrend MD, Schröter S, Stockle U, et al. (2018) Incidence and risk factors for postoperative Achilles tendon calcifications after percutaneous repair. *Arch Orthop Trauma Surg* 138:203-210
13. Ateschrang A, Riedmann S, Schröter S, Stöckle U (2013) Was man über Nahtanker zur Rekonstruktion der Rotatorenmanschett wissen sollte: wann welche Ankerkombination? *OP-Journal* 29:261-266
14. Ateschrang A, Schreiner AJ, Ahmad SS, Schröter S, Hirschmann MT, Korner D, et al. (2019) Improved results of ACL primary repair in one-part tears with intact synovial coverage. *Knee Surg Sports Traumatol Arthrosc* 27:37-43
15. Ateschrang A, Schröter S, Stöckle U (2015) Arthroskopie am oberen Sprunggelenk: Möglichkeiten und Grenzen. *OP-Journal* 30:112-118

16. Ateschrang AA, M.-D.; Stöckle, U.; Schröter, S.; Ahmad, S. (2018) Dynamische intraligamentäre Stabilisierung des vorderen Kreuzbandes. *Arthroskopie* 21:211-214
17. Dobele S, Gardner M, Schröter S, Hontzsch D, Stockle U, Freude T (2014) DLS 5.0--the biomechanical effects of dynamic locking screws. *PLoS One* 9:e91933
18. Döbele S, Schröter S, Hontzsch D, Stöckle U, Freude T (2012) Dynamic Locking Screw –DLS. *Op-Journal* 28:174-177
19. Ehnert S, Aspera-Werz RH, Freude T, Reumann MK, Ochs BG, Bahrs C, Schröter S, et al. (2016) Distinct Gene Expression Patterns Defining Human Osteoblasts' Response to BMP2 Treatment: Is the Therapeutic Success All a Matter of Timing? *Eur Surg Res* 57:197-210
20. Ehnert S, Aspera-Werz RH, Ihle C, Trost M, Zirn B, Flesch I, Schröter S, et al. (2019) Smoking Dependent Alterations in Bone Formation and Inflammation Represent Major Risk Factors for Complications Following Total Joint Arthroplasty. *J Clin Med* 8:
21. Ehnert S, Falldorf K, Fentz A, Ziegler P, Schröter S, Freude T, et al. (2015) Primary human osteoblasts with reduced alkaline phosphatase and matrix mineralization baseline capacity are responsive to extremely low frequency pulsed electromagnetic field exposure -clinical implication possible *Bone Reports*. *Bone reports* 3:48-56
22. Ehnert S, Schröter S, Aspera-Werz RH, Eisler W, Falldorf K, Ronniger M, et al. (2019) Translational Insights into Extremely Low Frequency Pulsed Electromagnetic Fields (ELF-PEMFs) for Bone Regeneration after Trauma and Orthopedic Surgery. *J Clin Med* 8:
23. Ehnert S, van Griensven M, Unger M, Scheffler H, Falldorf K, Fentz AK, Seeliger C, Schröter S, et al. (2018) Co-Culture with Human Osteoblasts and Exposure to Extremely Low Frequency Pulsed Electromagnetic Fields Improve Osteogenic Differentiation of Human Adipose-Derived Mesenchymal Stem Cells. *Int J Mol Sci* 19:
24. Fernandez Fernandez F, Eberhardt O, Schröter S, Wirth T, Ihle C (2019) Bilateral Tibial Tubercle Avulsion Fractures in Children -Clinical Results of a Rare Fracture. *Z Orthop Unfall*;10.1055/a-0979-2384
25. Floerkemeier S, Staubli AE, Schröter S, Goldhahn S, Lobenhoffer P (2014) Does obesity and nicotine abuse influence the outcome and complication rate after open-wedge high tibial osteotomy? A retrospective evaluation of five hundred and thirty three patients. *Int Orthop* 38:55-60
26. Floerkemeier S, Staubli AE, Schröter S, Goldhahn S, Lobenhoffer P (2013) Outcome after high tibial open-wedge osteotomy: a retrospective evaluation of 533 patients. *Knee Surg Sports Traumatol Arthrosc* 21:170-180
27. Freude T, Schröter S, Plecko M, Bahrs C, Martetschlaeger F, Kraus TM, et al. (2014) Dynamic-locking-screw (DLS)-leads to less secondary screw perforations in proximal humerus fractures. *BMC Musculoskelet Disord* 15:194
28. Freude T, Schröter S, Gonser CE, Stockle U, Acklin YP, Hontzsch D, et al. (2014) Controlled dynamic stability as the next step in „biologic plate osteosynthesis“ -a pilot prospective observational cohort study in 34 patients with distal tibia fractures. *Patient Saf Surg* 8:3
29. Freude T, Schröter S, Kraus TM, Hontzsch D, Stockle U, Dobele S (2013) [Dynamic locking screw 5.0--first clinical experience]. *Z Orthop Unfall* 151:284-290
30. Grunwald L, Angele P, Schröter S, Dickschas J, Harrer J, Hinterwimmer S, et al. (2019) Patients' expectations of osteotomies around the knee are high regarding activities of daily living. *Knee Surg Sports Traumatol Arthrosc* 27:3022-3031

31. Grunwald L, Dobele S, Hontzsch D, Slongo T, Stockle U, Freude T, Schröter S (2015) Callus massage after distraction osteogenesis using the concept of lengthening then dynamic plating. *Strategies Trauma Limb Reconstr* 10:129-135
32. Herbst M, Kuwashima U, Ahrend MD, Gueorguiev BG, Schröter S, Ihle C (2020) Health-Related Quality of Life -an Underestimated Factor to Evaluate the Treatment Success after Open Wedge HTO Surgery: Prospective 6-Years Follow-Up. *Z Orthop Unfall*;10.1055/a-1098-8894
33. Hill-Mundel K, Schlegl J, Biesalski HK, Ehnert S, Schröter S, Bahrs C, et al. (2019) Preoperative Ascorbic Acid Levels in Proximal Femur Fracture Patients Have No Postoperative Clinical Impact, While Ascorbic Acid Levels upon Discharge Have a Major Effect on Postoperative Outcome. *J Clin Med* 9:
34. Ihle C, Ahrend M, Grunwald L, Ateschrang A, Stockle U, Schröter S (2017) No change in patellar height following open wedge high tibial osteotomy using a novel femur-referenced measurement method. *Knee* 24:1118-1128
35. Ihle C, Ateschrang A, Albrecht D, Mueller J, Stockle U, Schröter S (2014) Occupational consequences after isolated reconstruction of the insufficient posterior cruciate ligament. *BMC Res Notes* 7:201
36. Ihle C, Ateschrang A, Grunwald L, Stockle U, Saier T, Schröter S (2016) Health-related quality of life and clinical outcomes following medial open wedge high tibial osteotomy: a prospective study. *BMC Musculoskelet Disord* 17:215
37. Ihle C, Maurer J, Ziegler P, Stockle U, Ateschrang A, Ahrend MD, Schröter S (2019) Sporting activity is reduced following medial reefing performed for patellar dislocation : A retrospective case series of 144 patients with a minimum follow-up of 24 months. *BMC Musculoskelet Disord* 20:34
38. Ihle C, Merk J, Schröter S, Ahrend M-D (2019) Rehabilitation nach kniegelenksnaher Umstellungsosteotomie. *Knie Journal* 1:36-44
39. Ihle C, Nakayama H, Gonser C, Stockle U, Dobele S, Schröter S, et al. (2018) [Suturing an Acute ACL Lesion with Dynamic Intraligamentary Stabilisation]. *Z Orthop Unfall* 156:100-102
40. Konstantinidis L, Grunewald H, Hauschild O, Schröter S, Hirschmuller A, Sudkamp NP, et al. (2015) Sliding of the load carrier in third-generation intramedullary nails for proximal femur fractures: an in vitro mechanical comparison study. *Proc InstMech Eng H* 229:110-114
41. Konstantinidis L, Papaioannou C, Hirschmuller A, Pavlidis T, Schröter S, Sudkamp NP, et al. (2013) Intramedullary nailing of trochanteric fractures: central or caudal positioning of the load carrier? A biomechanical comparative study on cadaver bones. *Injury* 44:784-790
42. Konstantinidis L, Papaioannou C, Hirschmuller A, Pavlidis T, Schröter S, Sudkamp NP, et al. (2012) Effects of muscle-equivalent forces on the biomechanical behavior of proximal femur fracture models: a pilot study on artificial bones. *Proc Inst Mech Eng H* 226:681-685
43. Konstantinidis L, Schmidt B, Bernstein A, Hirschmuller A, Schröter S, Sudkamp NP, et al. (2017) Plate fixation of periprosthetic femur fractures: What happens to the cement mantle? *Proc Inst Mech Eng H* 231:138-142
44. Korner D, Ateschrang A, Schröter S, Aurich M, Becher C, Walther M, et al. (2020) Concomitant ankle instability has a negative impact on the quality of life in patients with osteochondral lesions of the talus: data from the German Cartilage Registry (KnorpelRegister DGOU). *Knee Surg Sports Traumatol Arthrosc*;10.1007/s00167-020-05954-1
45. Korner D, Gueorguiev B, Niemeyer P, Bangert Y, Zinser W, Aurich M, Walther M, Becher C, Ateschrang A, Schröter S (2018) Correction to: Parameters influencing complaints and joint function in patients with osteochon-

- dral lesions of the ankle-an investigation based on data from the German Cartilage Registry (KnorpelRegister DGOU). Arch Orthop Trauma Surg 138:1333-1334
46. Korner D, Gueorguiev B, Niemeyer P, Bangert Y, Zinser W, Aurich M, Walther M, Becher C, Ateschrang A, Schröter S(2017) Parameters influencing complaints and joint function in patients with osteochondral lesions of the ankle-an investigation based on data from the German Cartilage Registry (KnorpelRegister DGOU). Arch Orthop Trauma Surg 137:367-373
47. Korner D, Kohler P, Schröter S, Naumann A, Walther M, Niemeyer P, et al. (2018) Pain in Osteochondral Lesions of the Ankle -an Investigation Based on Data from the German Cartilage Registry (KnorpelRegister DGOU). Z Orthop Unfall 156:160-167
48. Kraus TM, Abele C, Freude T, Ateschrang A, Stockle U, Stuby FM, Schröter S(2018) Duration of incapacity of work after tibial plateau fracture is affected by workintensity. BMC Musculoskelet Disord 19:281
49. Kraus TM, Freude T, Fiedler S, Schröter S, Stockle U, Ateschrang A (2015) Incapacity of work after arthroscopic Bankart repair. Arch Orthop Trauma Surg 135:1429-1436
50. Kraus TM, Imhoff AB, Ateschrang A, Stockle U, Schröter S(2015) [Clinical relevance of unloading in cartilage therapy of the knee--shoe insoles, knee braces or additional operative procedure?]. Z Orthop Unfall 153:75-79
51. Kuwashima U, Takeuchi R, Ishikawa H, Shioda M, Nakashima Y, Schröter S(2019) Comparison of torsional changes in the tibia following a lateral closed or medial open wedge high tibial osteotomy. Knee 26:374-381
52. Maiotti M, Massoni C, Russo R, Schröter S, Zanini A, Bianchedi D (2017) Arthroscopic Subscapularis Augmentation of Bankart Repair in Chronic Anterior Shoulder Instability With Bone Loss Less Than 25% and Capsular Deficiency: Clinical Multicenter Study. Arthroscopy 33:902-909
53. Maiotti M, Russo R, Zanini A, Schröter S, Massoni C, Bianchedi D (2016) Arthroscopic Bankart repair and subscapularis augmentation: an alternative technique treating anterior shoulder instability with bone loss. J Shoulder Elbow Surg 25:898-906
54. Nakayama H, Iseki T, Kanto R, Kambara S, Kanto M, Yoshiya S, Schröter S(2020) Physiologic kneejoint alignment and orientation can be restored by the minimally invasive double level osteotomy for osteoarthritic knees with severe varus deformity. Knee Surg Sports Traumatol Arthrosc 28:742-750
55. Nakayama H, Schröter S, Yamamoto C, Iseki T, Kanto R, Kurosaka K, et al. (2018) Large correction in opening wedge high tibial osteotomy with resultant joint-line obliquity induces excessive shear stress on the articular cartilage. Knee Surg Sports Traumatol Arthrosc 26:1873-1878
56. Notheisen TE, R.; Ruppert, E.; Gonser, C.; Schröter, S.; Fretschner, R. (2010) Ultraschallgesteuerte Regionalanästhesie in der Orthopädie/Traumatologie. Trauma und Berufskrankheit 12:128-133
57. Schaffler A, Dobele S, Stuby F, Maucher M, Schröter S, Stockle U, et al. (2014) [A new anatomical wing plate for osteoporotic acetabular fractures: biomechanical testing and first clinical experience]. Z Orthop Unfall 152:26-32
58. Schröter S, Stöckle U, Ateschrang A (2016) Surgical methods to unload the knee [Operative Methoden zur Entlastung des Kniegelenks]. OUP 10:568-575
59. Schröter S, Weise K, Badke A (2009) [Traumatic lumbosacral dislocation -an underrated injury]. Z Orthop Unfall 147:231-235
60. Schröter S(2013) Critical comments and questions to thepublication „Getgood A, Collins B, Slynarski K, Kurowska E, Parker D, Engebretsen L, MacDonald PB, Litchfield R 2011 Short-term safety and efficacy of a novel

high tibial osteotomy system: a case controlled study. *Knee Surg Sports Traumatol Arthrosc*. *Knee Surg Sports Traumatol Arthrosc* 21:270-271

61. Schröter S, Ateschrang A (2014) Critical comments and questions regarding the article „serial assessment of weight-bearing lower extremity alignment radiographs after open-wedge high tibial osteotomy“. *Arthroscopy* 30:1050-1051

62. Schröter S, Ateschrang A, Flesch I, Stockle U, Freude T (2016) First mid-term results after cancellous allograft vitalized with autologous bone marrow for infected femoral non-union. *Wien Klin Wochenschr* 128:827-836

63. Schröter S, Ateschrang A, Ihle C, Stockle U, Konstantinidis L, Dobeles S (2014) [Lateral hinge fractures in open wedge high tibial osteotomy]. *Orthopade* 43:1000-1007

64. Schröter S, Ateschrang A, Lowe W, Nakayama H, Stockle U, Ihle C (2017) Early full weight-bearing versus 6-week partial weight-bearing after open wedge high tibial osteotomy leads to earlier improvement of the clinical results: a prospective, randomised evaluation. *Knee Surg Sports Traumatol Arthrosc* 25:325-332

65. Schröter S, Elson DW, Ateschrang A, Ihle C, Stockle U, Dickschas J, et al. (2017) Lower Limb Deformity Analysis and the Planning of an Osteotomy. *J Knee Surg* 30:393-408

66. Schröter S, Freude T, Kopp MM, Konstantinidis L, Dobeles S, Stockle U, et al. (2015) Smoking and unstable hinge fractures cause delayed gap filling irrespective of early weight bearing after open wedge osteotomy. *Arthroscopy* 31:254-265

67. Schröter S, Gonser C, Ateschrang A, Albrecht D, Weise K (2010) Experience with computer assisted navigation in high tibial osteotomy [Hohe Tibiakopfumstellungsosteotomie]. *Trauma und Berufskrankheit* 12:95-100

68. Schröter S, Gonser CE, Konstantinidis L, Helwig P, Albrecht D (2011) High Complication Rate After Biplanar Open Wedge High Tibial Osteotomy Stabilized With a New Spacer Plate (Position HTO Plate) Without Bone Substitute. *Arthroscopy: The Journal of Arthroscopic & Related Surgery* 27:644-652

69. Schröter S, Gunzel J, Freude T, Ateschrang A, Stockle U, Albrecht D (2012) [Precision in the planning of open wedge HTO]. *Z Orthop Unfall* 150:368-373

70. Schröter S, Hoffmann T, Dobeles S, Welke B, Hurschler C, Schwarze M, et al. (2018) Biomechanical properties following open wedge high tibial osteotomy: Platefixator combined with dynamic locking screws versus standard locking screws. *Clin Biomech (Bristol, Avon)* 60:108-114

71. Schröter S, Ihle C, Elson DW, Dobeles S, Stockle U, Ateschrang A (2016) Surgical accuracy in high tibial osteotomy: coronal equivalence of computer navigation and gap measurement. *Knee Surg Sports Traumatol Arthrosc* 24:3410-3417

72. Schröter S, Ihle C, Mueller J, Lobenhoffer P, Stockle U, van Heerwaarden R (2013) Digital planning of high tibial osteotomy. Interrater reliability by using two different software. *Knee Surg Sports Traumatol Arthrosc* 21:189-196

73. Schröter S, Konstantinidis L, Kraus TM, Stuby F, Stockle U, Freude T (2013) Open wedge re-correction high tibial osteotomy in an elderly patient with a varus angulated non-united dome-shaped high tibial osteotomy. *J Zhejiang Univ Sci B* 14:1054-1058

74. Schröter S, Kramer M, Welke B, Hurschler C, Russo R, Herbst M, et al. (2016) The effect of the arthroscopic augmentation of the subscapularis tendon on shoulder instability and range of motion: A biomechanical study. *Clin Biomech (Bristol, Avon)* 38:75-83

75. Schröter S, Lobenhoffer P, Mueller J, Ihle C, Stockle U, Albrecht D (2012) [Changes of patella position after closed and open wedge high tibial osteotomy: review of the literature]. Orthopade 41:186, 188-194
76. Schröter S, Lutz O, Gonser CE, Ateschrang A, Badke A, Albrecht D (2011) [Postarthroscopic glenohumeral chondrolysis--are there any causal factors? Case report]. Z Orthop Unfall 149:688-693
77. Schröter S, Mueller J, van Heerwaarden R, Lobenhoffer P, Stockle U, Albrecht D (2013) Return to work and clinical outcome after open wedge HTO. Knee Surg Sports Traumatol Arthrosc 21:213-219
78. Schröter S, Nakayama H, Ihle C, Ateschrang A, Maiotti M, Harrer J, et al. (2019) Torsional Osteotomy. J Knee Surg;10.1055/s-0039-1678677
79. Schröter S, Nakayama H, Yoshiya S, Stockle U, Ateschrang A, Gruhn J (2019) Development of the double level osteotomy in severe varus osteoarthritis showed good outcome by preventing oblique joint line. Arch Orthop Trauma Surg 139:519-527
80. Ziegler P, Nussler AK, Wilbrand B, Falldorf K, Springer F, Fentz AK, Eschenburg G, Ziegler A, Stöckle U, Maurer E, Ateschrang A, Schröter S, Ehnert S(2019) Pulsed Electromagnetic Field Therapy Improves Osseous Consolidation after High Tibial Osteotomy in Elderly Patients-A Randomized, Placebo-Controlled, Double-Blind Trial. J Clin Med 8: